Cornwallis Academy
Ambitions for All

SMILE
SUCCESSFUL | MOTIVATED | INSPIRED | LEARNING | EXCITED & ENGAGED

Year 13 A Level Phyics

3.6 Further mechanics and thermal physics

Learning Objectives

By the end of this unit, students should be able to:

- **Circular Motion:** Define angular velocity (omega = v/r = 2pi f). Calculate centripetal acceleration (a = v2/r = omega2 r) and centripetal force (F = mv2/r = m omega2 r), applying these to vertical loops, amusement park rides, and banked tracks.
- **Simple Harmonic Motion (SHM):** Define SHM by the characteristic equation a = -omega2 x. Analyze position, velocity, and acceleration graphs against time, identifying phase relationships.
- **SHM Systems:** Apply mass-spring system (T = 2pi sqrt m/k) and simple pendulum (T = 2pi sqrt l/g) time period formulas to solve problems and determine experimental constants.
- **Damping and Resonance:** Distinguish between light, heavy, and critical damping. Describe free and forced oscillations, and explain the conditions required for resonance, including how damping affects the resonance peak.
- **Thermal Energy Transfer:** Define specific heat capacity (Q = mc Delta theta) and specific latent heat (Q = mL). Interpret heating and cooling curves.
- **Ideal Gases:** State and apply the ideal gas laws (Boyle's, Charles's, and Pressure laws). Use the ideal gas equations pV = nRT and pV = NkT.
- **Kinetic Theory Model:** Explain the macroscopic properties of a gas (pressure) using a microscopic kinetic model. Understand the derivation of pV = 1/3 Nm c2 and relate average molecular kinetic energy to absolute temperature (Ek = 3/2 kT).

What previous knowledge is required?

Students will build heavily upon core concepts from Year 12 (AS) Physics and Mathematics:

- **From Section 3.4 (Mechanics):** Newton's laws of motion, work, energy conservation, and resolving vectors into components.
- **From Section 3.2 (Waves):** Fundamental terminology such as frequency (f), period (T), amplitude, and phase difference.
- **From Mathematics:** Radian measure, trigonometric graphs (sin and cos), and a conceptual or calculus-based understanding of gradients (velocity as the gradient of displacement, acceleration as the gradient of velocity)

How will this term's content help with future learning?

- **Section 3.7 (Fields and their Consequences):** The mathematics of circular motion is used continuously to analyze satellite orbits in gravitational fields and the circular deflection of charged particles in magnetic fields.
- **Section 3.8 (Nuclear Physics):** The kinetic theory of gases and thermal energy concepts are essential for understanding the conditions required for nuclear fusion in stars and the cooling mechanisms within nuclear reactors.
- **Astrophysics Option:** Thermal physics, ideal gases, and molecular kinetic energies form the exact physics required to model stellar classification, blackbody radiation, and core stellar dynamics.

Common Misconceptions

- **The Centripetal Force Concept:** Students frequently treat centripetal force as an independent, additional force acting on an object, rather than identifying it as the *resultant* of existing forces (e.g., gravity, tension, or friction) directed toward the center of rotation.

- **SHM Acceleration Direction:** Students often mistakenly believe that acceleration points in the direction of motion. They must be constantly reminded that in SHM, acceleration is *always* directed toward the equilibrium position, meaning it opposes displacement (a propto –x).
- **Resonance Peak and Damping:** Students often fail to realize that increasing damping not only lowers the maximum amplitude at resonance but also shifts the resonant frequency slightly to a *lower* frequency.
- **Temperature Scales:** In gas law calculations, students frequently forget to convert temperatures from degrees Celsius (°C) to Kelvin (K), which completely invalidates calculations using pV = nRT.
- **Gas Law Assumptions:** Students treat the ideal gas equations as flawless, absolute laws. They struggle to articulate the microscopic assumptions of the kinetic theory model (e.g., that intermolecular forces are negligible except during collisions, and the volume of gas molecules is negligible compared to the container volume).

National Curriculum / Specification

- **3.6.1 Further Mechanics:**
  - 3.6.1.1 Uniform circular motion (Angular speed, centripetal acceleration, and force equations).
  - 3.6.1.2 Simple harmonic motion (SHM definitions, displacement, velocity, and acceleration equations).
  - 3.6.1.3 Simple harmonic systems (Mass-spring system, simple pendulum).
  - 3.6.1.4 Forced vibrations and resonance (Damping, critical damping, resonance applications).
- **3.6.2 Thermal Physics:**
  - 3.6.2.1 Thermal energy transfer (Internal energy, specific heat capacity, specific latent heat).
  - 3.6.2.2 Ideal gases (Boyle's law, Charles's law, ideal gas equation, Avogadro and Boltzmann constants).
  - 3.6.2.3 Molecular kinetic theory model (Pressure derivation, root-mean-square speed, molecular kinetic energy relation).

Required Practicals Covered in this Section:

- **Required Practical 7:** Investigation into simple harmonic motion using a mass-spring system and a simple pendulum.
- **Required Practical 8:** Investigation of Boyle's law and Charles's law to check the relationships between pressure, volume, and temperature of a gas.

3.7 Fields and their consequences

Learning Objectives

By the end of this unit, students should be able to:

- **Fields Concept:** Define a force field and distinguish between gravitational, electric, and magnetic fields, recognizing similarities and differences in their governing inverse-square laws.
- **Gravitational Fields:** Apply Newton’s law of gravitation (F = G M m / r2). Define and calculate gravitational field strength (g), gravitational potential (V), and gravitational potential energy. Analyze planetary and satellite orbits, including geostationary orbits and escape velocity.
- **Electric Fields:** Apply Coulomb’s law (F =1\4pi varepsilon0 Q1 Q2/r2). Define electric field strength (E) and electric potential (V) for both uniform and radial fields. Analyze the trajectories of moving charges in uniform electric fields.
- **Capacitance:** Define capacitance (C = Q/V). Calculate the energy stored in a capacitor. Analyze the exponential charging and discharging rates of capacitors through resistors, utilizing the time constant (tau = RC).

- **Magnetic Fields:** Determine the force on a current-carrying wire (F = BIl) and a moving charge (F = Bqv) in a magnetic field using Fleming's Left-Hand Rule. Explain the circular trajectory of a charged particle in a uniform magnetic field.
- **Electromagnetic Induction:** Apply Faraday's law and Lenz's law to explain induced electromotive force (emf). Calculate magnetic flux (Phi = BA) and flux linkage N Phi). Analyze alternating current (AC) generators and the construction/efficiency of transformers.

What previous knowledge is required?

- **From Section 3.3 (Electricity):** Current, potential difference, resistance, and exponential relationships.
- **From Section 3.4 & 3.6 (Mechanics / Circular Motion):** Newton's laws of motion, work done, kinetic and potential energy conservation, angular speed, and centripetal acceleration (a = v2/r).
- **From Mathematics:** Working fluently with natural logarithms (ln) and exponentials (e), vector resolution, and inverse-square proportionality.

How will this term's content help with future learning?

- **Section 3.8 (Nuclear Physics):** The concepts of electric potential energy and Coulombic repulsion are vital for understanding the Rutherford alpha particle scattering experiment, distance of closest approach, and the conditions required to overcome repulsion for nuclear fusion.
- **Medical Physics / Turning Points Options:** The behavior of charged particles in magnetic and electric fields directly underpins the mechanics of particle accelerators (cyclotrons, synchrotrons), cathode ray tubes, and medical imaging devices like MRI scanners.

Common Misconceptions

- **Field Lines vs. Particle Paths:** Students often think electric or gravitational field lines show the exact path a moving object will take. They must be corrected to understand that field lines show the direction of the *force* acting on a stationary test mass/charge at that specific point.
- **Gravitational vs. Electric Potential:** Students often confuse the two potentials. Critically, they forget that gravitational potential is *always negative* (as it is an attractive field relative to infinity), whereas electric potential can be positive or negative depending on the sign of the charge creating the field.
- **Capacitor Discharge Behavior:** Students frequently believe that a capacitor discharges at a constant linear rate. They struggle to visualize the exponential decay, where the rate of discharge drops as the remaining charge (and thus driving voltage) decreases.
- **Lenz's Law Signs:** When applying Faraday's and Lenz's laws (varepsilon = -Delta (N Phi)/ Delta t), students often ignore the negative sign. They treat it as a trivial detail rather than the mathematical representation of the conservation of energy (the induced current must oppose the change creating it).
- **Magnetic Force Direction:** Students often struggle to apply Fleming's Left-Hand Rule to *negative* charges (like electrons). They fail to reverse their second finger (current), which represents the flow of conventional *positive* charge.

National Curriculum / Specification

- **3.7.1 Fields:** Concept of a field and comparison of field types.
- **3.7.2 Gravitational Fields:** Newton's law, radial/uniform fields, gravitational potential, satellite orbits.
- **3.7.3 Electric Fields:** Coulomb's law, field strength E, electric potential, path of charged particles.
- **3.7.4 Capacitance:** Capacitors in series/parallel, energy storage, charging and discharging equations.
- **3.7.5 Magnetic Fields:** Magnetic flux density B, force on moving charges, cyclotrons, Faraday's and Lenz's laws, AC generators, and transformers.

Required Practicals Covered in this Section:

- **Required Practical 9:** Investigation of the charge and discharge of a capacitor. An analysis of data to determine the time constant (tau = RC) using log-linear graphing.
- **Required Practical 10:** Investigate how the force on a current-carrying conductor in a magnetic field varies with current, length of wire, and magnetic flux density (using a top-pan balance).

3.8 Nuclear physics

Learning Objectives

- **Rutherford Scattering:** Describe the Rutherford alpha particle scattering experiment and explain how the results provided evidence for a nuclear atom. Calculate the distance of closest approach to estimate nuclear radius.
- **Nuclear Radius and Density:** State and apply the nuclear radius equation (R = R0 A1/3). Explain the experimental method of high-energy electron diffraction to determine nuclear radius, and show that nuclear density is constant.
- **Radioactive Decay:** Compare the properties of alpha, beta-, beta+, and gamma radiation. State and apply the exponential decay equations (N = N0e -lambda t and A = A0e -lambda t, and define decay constant (lambda) and half-life (T1/2 = ln 2 / lambda).
- **Nuclear Stability:** Interpret N against Z graphs to identify modes of decay (alpha, beta-, beta+, and electron capture) for unstable nuclei.
- **Mass Defect and Binding Energy:** Define mass defect and binding energy. Use Einstein's equation (Delta E = Delta m c2) to calculate binding energy and binding energy per nucleon.
- **Fission and Fusion:** Interpret the graph of binding energy per nucleon against nucleon number (A) to explain the processes of nuclear fission and nuclear fusion.
- **Nuclear Reactor Physics:** Explain the function of fuel rods, moderators (thermal neutrons), control rods, and coolants in a thermal nuclear reactor. Discuss the environmental impacts of nuclear waste handling and disposal.

What previous knowledge is required?

- **From Section 3.1 (Particles and Radiation):** Basic atomic structures, isotopes, fundamental conservation laws (baryon and lepton numbers), and the properties of alpha and beta decay.
- **From Section 3.4 (Mechanics):** Conservation of momentum and kinetic energy during collisions and recoil events.
- **From Section 3.7 (Fields):** Coulomb's law and electric potential energy (Ep = 1 / 4pi varepsilon0 Q1 Q2 /r), which is vital for calculating the distance of closest approach.

How will this term's content help with future learning?

- **Synoptic Linkage (Papers 1, 2, and 3):** This unit relies heavily on data interpretation, logarithms, and modeling exponential decay, which are core skills tested across all three final written exam papers.
- **Astrophysics Option:** Binding energy curves, the proton-proton chain, and the conditions required to overcome Coulombic repulsion for nuclear fusion directly underpin the stellar evolution models studied in the Astrophysics module.
- **Turning Points in Physics Option:** The wave-particle duality concepts used in electron diffraction to measure the nuclear radius directly link to the "Turning Points" exploration of the dual nature of matter.

Common Misconceptions

- **losest Approach vs. Diffraction:** Students often think the distance of closest approach gives the *exact* radius of the nucleus. They struggle to realize it only yields an *upper limit* because the alpha particle is

stopped by electrostatic repulsion before touching the nucleons, whereas electron diffraction provides a much more precise measurement.

- **Nuclear Density Dependency:** Students frequently assume that heavier nuclei (higher A) must be vastly denser than lighter ones. They fail to mathematically deduce from R = R0 A1/3 that volume (V propto R3) is proportional to mass (M propto A), making nuclear density constant across all elements.
- **Activity vs. Count Rate:** Students often treat the actual activity (A) of a sample as identical to the measured count rate on a Geiger-Müller tube, forgetting to correct for background radiation or the geometric capture efficiency of the detector.
- **Binding Energy vs. Total Energy:** When calculating mass defect, students often subtract the mass of the nucleus from the constituent nucleons but get confused about the sign of the energy, struggling to realize that binding energy is the energy *released* when a nucleus forms, or the energy *required* to separate it.
- **Moderator vs. Control Rods:** Students continuously mix up the roles of reactor components. They often state that a moderator "slows down the reaction," when in fact it slows down the *neutrons* (turning them into thermal neutrons) to actually *increase* the probability of further fission. Control rods are what control or slow down the rate of the reaction by absorbing neutrons.

National Curriculum / Specification

- **3.8.1 Radioactivity:**
  - 3.8.1.1 Rutherford scattering (Evidence for the nucleus, alpha scattering observations).
  - 3.8.1.2 alpha, beta and gamma radiation (Properties, inverse-square law for gamma radiation).
  - 3.8.1.3 Radioactive decay (Decay constant lambda, half-life, exponential equations).
  - 3.8.1.4 Nuclear instability (Graph of N against Z, decay modes, excited states and gamma emission).
  - 3.8.1.5 Nuclear radius (High-energy electron diffraction, R = R0 A1/3, constant density calculation).
  - 3.8.1.6 Mass and energy (E=mc2, mass defect, binding energy, binding energy per nucleon curve).
  - 3.8.1.7 Induced fission and safety (Thermal nuclear reactors, fuel, moderator, control rods, coolant, safety features, high/low-level radioactive waste disposal).

Required Practicals Covered in this Section:

- **Required Practical 11:** Investigation of the inverse-square law for gamma radiation using a Geiger-Müller tube and a gamma source (such as Cobalt-60), incorporating corrections for background radiation.

3.12 Turning Points Optional Unit

Learning Objectives

By the end of this unit, students should be able to:

- **The Discovery of the Electron:** Describe cathode rays and explain Thomson’s experiment to determine the specific charge of the electron (e/m). Explain Millikan’s oil drop experiment and how it proved the quantization of electric charge.
- **Wave-Particle Duality:** Explain Newton’s corpuscular theory of light versus Huygens’ wave theory, and describe how Foucault's measurement of the speed of light in water resolved the debate. Explain the significance of Maxwell’s electromagnetic theory and Hertz’s discovery of radio waves.
- **The Blackbody Crisis and Quantum Mechanics:** Explain the "ultraviolet catastrophe" and Planck’s quantization hypothesis. Revisit the photoelectric effect and explain how Einstein’s photon theory revolutionized the understanding of light.
- **Electron Microscopes:** Compare the resolution limits of optical microscopes to electron microscopes using de Broglie’s wavelength equation, analyzing transmission (TEM) and scanning (SEM) electron microscope mechanics.

- **Special Relativity:** Explain the significance of the Michelson-Morley experiment and its null result regarding the "absolute ether". State Einstein's two postulates of special relativity.
- **Relativistic Consequences:** Calculate and explain time dilation (t = t0 / sqrt (1 - v2/c2)), length contraction (l = l0 sqrt (1 - v2/c2)), and relativistic mass/energy equivalence (E = mc2 = m0c2 / sqrt (1 - v2/c2)). Interpret experimental evidence for relativity, such as muon decay.

What previous knowledge is required?

- **From Section 3.1 & 3.2 (Particles & Waves):** Basic wave mechanics, interference patterns, photons (E=hf), and the preliminary concept of wave-particle duality.
- **From Section 3.4 (Mechanics):** Work done, kinetic energy formulas, and basic classical momentum (p=mv).
- **From Section 3.7 (Fields):** The motion of charged particles in uniform electric fields (F=EQ) and magnetic fields (F=Bqv), which is foundational for understanding cathode ray tubes, Thomson's crossed fields, and Millikan's balanced forces.

How will this term's content help with future learning?

Because "Turning Points" is traditionally taught at the absolute end of Year 13, it serves less as a bridge to later chapters and more as a bridge to **Higher Education (University Level Physics)**:

- **Preparation for Quantum Mechanics:** The progression from classical thermodynamics (blackbody radiation) to Planck's quanta and de Broglie matter waves prepares students for undergraduate quantum physics and operators.
- **Preparation for Modern Astrophysics:** Relativistic mass scaling, time dilation, and space-time transformations are fundamental for students going on to study cosmology, general relativity, or high-energy particle physics.
- **Synoptic Exam Execution:** The unit heavily reinforces field mechanics and particle physics calculations, meaning it acts as excellent active revision for the highly weighted **Paper 1 and Paper 2 core questions**.

Common Misconceptions

- **Specific Charge vs. Individual Charge:** Students often mistake J.J. Thomson's determination of the electron's *specific charge* (e/m) as the discovery of the absolute charge of the electron (e), forgetting that it was Millikan who isolated the individual charge value later on.
- **Millikan's Terminal Velocity:** In Millikan's experiment, students frequently assume the electric field is turned on during the entire drop. They fail to understand that the field is initially *off* so the drop can reach a terminal velocity under gravity/drag alone, which is vital to calculate the drop's radius and mass.
- **Proper Time (t0) and Proper Length (l0):** This is the most prevalent error in relativity. Students struggle to identify which frame of reference measures the "proper" metric. They must be explicitly coached that proper time (t0) is *always* measured by an observer who is stationary relative to the event being timed.
- **"Mass Increasing" Intuition:** Students often conceptually struggle with why an object cannot accelerate past the speed of light (c). They fail to see that as v --> c , the relativistic energy required to yield a minor increase in velocity approaches infinity, meaning the work input changes the relativistic mass/momentum rather than increasing speed linearly.
- **The Ether Null Result:** Students often think the Michelson-Morley experiment was a failure because it returned a "null result." They need to learn that in physics, a precise null result can be a profound success, as it successfully disproved the existence of the luminiferous ether.

National Curriculum / Specification

- **3.12.1 The Discovery of the Electron:**
  - 3.12.1.1 Cathode rays (Production, thermionic emission, specific charge e/m determination).

- 3.12.1.2 Millikan’s experiment (Quantization of charge, viscous drag, Stokes' law).

- **3.12.2 Wave-Particle Duality:**
  - 3.12.2.1 Newton’s corpuscular theory vs. Huygens' wave theory (Diffraction, speed of light in media).
  - 3.12.2.2 Significance of Maxwell’s waves & Hertz’s discoveries (EM wave propagation, radio detection).
  - 3.12.2.3 Clean-up of the Quantum crisis (Blackbody curves, photo-electricity validation).
  - 3.12.2.4 Electron microscopes (De Broglie applied to TEM and SEM systems).
- **3.12.3 Special Relativity:**
  - 3.12.3.1 The Michelson-Morley experiment (Interferometer principles, speed of light invariability).
  - 3.12.3.2 Einstein’s postulates (Inertial frames of reference, consistency of c).
  - 3.12.3.3 Relativistic consequences (Time dilation, length contraction, cosmic muon validation).
  - 3.12.3.4 Mass and energy (Relativistic mass equation, kinetic energy modifications at high velocities).